

## 443

**Reduced-Intensity Stem Cell Allografting for PNH**

**Patients in the Eculizumab Era: The Mexican Experience**  
 Alejandro Schcolnik-Cabrera<sup>1</sup>, Nancy Labastida-Mercado<sup>2,3</sup>,  
 Samantha L. Galindo-Becerra<sup>2,3</sup>, David Gomez-Almaguer<sup>4</sup>,  
 Miguel A. Herrera-Rojas<sup>4</sup>, **Guillermo J. Ruiz-Delgado**<sup>2,3,5</sup>,  
 Guillermo J. Ruiz-Arguelles<sup>5,6,7</sup>, <sup>1</sup>Universidad Nacional  
 Autónoma de México, Mexico City, Mexico; <sup>2</sup>Centro de  
 Hematología y Medicina Interna, Clínica Ruiz de Puebla,  
 Puebla, Mexico; <sup>3</sup>Universidad Popular Autónoma del Estado  
 de Puebla, Puebla, Mexico; <sup>4</sup>Hematology Service, University  
 Hospital of Monterrey, Monterrey, Mexico; <sup>5</sup>Laboratorios  
 Clínicos de Puebla, Clínica Ruiz, Puebla, Mexico; <sup>6</sup>Centro de  
 Hematología y Medicina Interna, Clínica Ruiz, Puebla,  
 Mexico; <sup>7</sup>Universidad de las Américas Puebla, Puebla,  
 Mexico

**Background:** Paroxysmal nocturnal haemoglobinuria (PNH) presents as two major entities: the classical form, predominantly hemolytic, and a secondary type with marrow failure and resultant aplastic anaemia (AA-PNH). Currently, the treatment of choice of the hemolytic variant is eculizumab; however, the most frequent form of PNH in México is AA-PNH.

**Patients and Methods:** Six consecutive AA-PNH patients with HLA-identical siblings were allografted in two institutions in México, employing a reduced-intensity conditioning regimen for stem cell transplantation (RIST) conducted on an outpatient basis.


**Results:** Median age of the patients was 37 years (range 25–48). The patients were given a median of  $5.4 \times 10^6/\text{kg}$  allogeneic CD34(+) cells, using 1–3 apheresis procedures. Median time to achieve above  $0.5 \times 10^9/\text{l}$  granulocytes was 21 days, whereas median time to achieve above  $20 \times 10^9/\text{l}$  platelets was 17 days. Five patients are alive for 330–3150 days (median 1437) after the allograft. The 3150-day overall survival is 83.3%, whereas median survival has not been reached, being above 3150 days.

**Conclusion:** We have shown that hypoplastic PNH patients can be allografted safely using RIST and that the long-term results are adequate, the cost–benefit ratio of this treatment being reasonable. Additional studies are needed to confirm the usefulness of RIST in the treatment of AA-PNH.

## 444

**Ibrutinib Treatment of Relapsed CLL Following Allogeneic Transplantation: Sustained Disease Response and Promising Donor Immune Modulation**

**Christine E. Ryan**<sup>1</sup>, Aaron C. Logan<sup>2</sup>, Andrew Rezvani<sup>1</sup>,  
 Manali Kamdar<sup>1</sup>, Hideki Nakasone<sup>1</sup>, Bitu Sahaf<sup>1</sup>,  
 Joanne Otani<sup>1</sup>, Katherine A. Kong<sup>3</sup>, Mark Klinger<sup>3</sup>,  
 Malek Faham<sup>3</sup>, Steven Coutre<sup>4</sup>, David B. Miklos<sup>1</sup>. <sup>1</sup>Division  
 of Blood and Marrow Transplantation, Stanford University  
 School of Medicine, Stanford, CA; <sup>2</sup>Helen Diller Family  
 Comprehensive Cancer Center, University of California San  
 Francisco, San Francisco, CA; <sup>3</sup>Sequentia, Inc., South San  
 Francisco, CA; <sup>4</sup>Division of Hematology, Stanford University  
 School of Medicine, Stanford, CA



**Figure 1.** (A) Percent reduction in LN size, reported as sum of the product of LN diameters (SPD), for 4 patients following ibrutinib initiation. (B) CLL MRD (as percent of WBCs) and blood donor CD3 chimerism for SPN3975 and (C) SPN3431. (D) B cells (excluding the CLL clone) as percent of total PBMC for SPN3975. (E) Total IgH molecules and unique IgH clone counts for SPN3975 at time points (D=day) post allo-HCT.